

Variety Description

Variety

TRERANO

| | |
|------------------------------|--------------------|
| Species | Cocksfoot |
| Botanical name | Dactylis glomerata |
| Ploidy | 2 |
| Seeding rate | 20–25 kg/ha |
| Distance between rows | as cereals |
| Sowing period | April to August |
| Sowing depth | 1–2 cm |

Agronomic figures*:

| | |
|---------------------------|---|
| Panicle formation | 2 |
| Development after sowing | 7 |
| Tendency to winterkilling | 4 |
| Tendency to lodging | 6 |
| Susceptibility to rust | 6 |
| Sward density | 5 |
| DM-yield total | 6 |
| DM-yield first cut | 7 |
| DM-yield in aftermath | 4 |



Clarification of figures*:

1: very early, very low / 5: medium / 9: very late, very high

* **Source:** Bundessortenamt [Federal Plant Variety Office] 2018, state variety tests

Variety description

TRERANO is an early cocksfoot variety with excellent yield qualities. It produces high yields from the first cut and forms abundant mass in the early development stages. All these qualities result in excellent stand and weed suppression. In addition, the variety is also very winter-hardy.

Most important characteristics

Early maturity group
High mass formation in the early development
Good winter-hardiness
High yield in first cut
High overall yield

Usage

Cocksfoot or orchard grass is considered ideal for dry places and is resistant to cold. It can tolerate intensive fertilisation level and cropping, as well as grazing. The variety can be used for 4–6 grazings and be mowed 3–4 times.

